An overdose happens when someone takes too much of an opioid.

A person who has overdosed will be unable to wake up or talk to you.

Their breathing may be slow and shallow.

> They may not be breathing at all.



FindNaloxoneNowKY.org

Give Naloxone!

Naloxone reverses overdose. Act quickly!

Peel Peel back the

foil and remove

Place

Gently insert the nozzle into the nasal spray, their nostril.

Press the plunger all the way in until it clicks.

Press



Naloxone nasal spray takes 2-3 minutes to work! Wait at least 2 minutes for a response before giving more. CALL 911! Give rescue breaths while you wait.

An overdose happens when someone takes too much of an opioid.

A person who has overdosed will be unable to wake up or talk to you.

Their breathing may be slow and shallow.

> They may not be breathing at all.



Find Naloxone NowKY

FindNaloxoneNowKY.org

Give Naloxone

Naloxone reverses overdose. Act quickly!

Place Peel

Peel back the foil and remove the nasal spray. their nostril.

Gently Insert the nozzle into Press the plunger all the way in until it clicks.

Press



Naloxone nasal spray takes 2-3 minutes to work! Vait at least 2 minutes for a response before giving more CALL 911! Give rescue breaths while you wait.

An overdose happens when someone takes too much of an opioid.

A person who has overdosed will be unable to wake up or talk to you

Their breathing may be slow and shallow.

They may not be breathing at all.



Naloxone Nowice

indNaloxoneNowKY.org

Give Naloxone

Naloxone reverses overdose. Act quickly!

Place Peel

Peel back the foil and remove the nozzle into the nasal spray.

Gently Insert their nostriL

Press the plunger all the way in until it clicks.

Press

Press



Naloxone nasal spray takes 2-3 minutes to work! Vait at least 2 minutes for a response before giving mo CALL 911! Give rescue breaths while you wait.

An overdose happens when someone takes too much of an opioid.

A person who has overdosed will be unable to wake up or talk to you.

Their breathing may be slow and shallow.

> They may not be breathing at all.



FindNaloxoneNowKY.org

Give Naloxone

Naloxone reverses overdose. Act quickly!

<u>Place</u>

Peel

Peel back the Gently Insert foil and remove the nozzle into the nasal spray.



Press the plunger all the way in until it clicks.

Press



Naloxone nasal spray takes 2-3 minutes to work! it at least 2 minutes for a response before giving more CALL 911! Give rescue breaths while you wait.

An overdose happens when someone takes too much of an opioid.

A person who has overdosed will be unable to wake up or talk to you.

Their breathing may be slow and shallow

> They may not be breathing at all.



Find Naloxone

FindNaloxoneNowKY.org An overdose happens

when someone takes too

much of an opioid.

A person who has

overdosed will be unable

to wake up or talk to you.

Their breathing may be

slow and shallow.

They may not be

breathing at all.

Give Naloxone!

Naloxone reverses overdose. Act quickly! Place

Peel

Peel back the

Gently Insert foil and remove the nozzle into the nasal spray. their nostril.





An overdose happens hen someone takes too much of an opioid.

A person who has overdosed will be unable to wake up or talk to you.

Their breathing may be slow and shallow.

> They may not be breathing at all.



Find Naloxone NowKY

FindNaloxoneNowKY.org

Give Naloxone!

Naloxone reverses overdose. Act quickly!

Peel

Peel back the foil and remove the nozzle into

Gently Insert

Place

the nasal spray. their nostril.

Press the plunger all the way in until it clicks.

Press



Naloxone nasal spray takes 2-3 minutes to work! Vait at least 2 minutes for a response before giving more CALL 911! Give rescue breaths while you wait.

Give Naloxone!

CALL 911! Give rescue breaths while you wait.

Naloxone reverses overdose. Act quickly!

Peel

Peel back the foil and remove the nasal spray. their nostril.

Place Gently Insert the nozzle into

Press Press the plunger all the way in until



Naloxone nasal spray takes 2-3 minutes to work! Vait at least 2 minutes for a response before giving more. CALL 911! Give rescue breaths while you wait. -

An overdose happens when someone takes too much of an opioid.

A person who has overdosed will be unable to wake up or talk to you.

Their breathing may be slow and shallow.

> They may not be breathing at all.



FindNaloxoneNowKY.org

Give Naloxone

Naloxone reverses overdose. Act quickly!

Peel

Peel back the foil and remove

Place

Gently Insert the nasal spray. their nostril.

the nozzle into

Press Press the plunger all the way in until



Naloxone nasal spray takes 2-3 minutes to work! Wait at least 2 minutes for a response before giving more CALL 911! Give rescue breaths while you wait.

indNaloxoneNowKY.org An overdose happens when someone takes too

much of an opioid. A person who has overdosed will be unable to wake up or talk to you.

Their breathing may be slow and shallow They may not be



ndNaloxoneNowKY.or

Give Naloxone!

Naloxone reverses overdose. Act quickly!

Peel Peel back the

Place Gently Insert

foil and remove the nozzle into

Press Press the plunger all the way in until it clicks.



Naloxone nasal spray takes 2-3 minutes to work! Vait at least 2 minutes for a response before giving more CALL 911! Give rescue breaths while you wait.

An overdose happens when someone takes too much of an opioid.

A person who has overdosed will be unable to wake up or talk to you.

Their breathing may be slow and shallow. They may not be

breathing at all.



FindNaloxoneNowKY.org

Give Naloxone!

Naloxone reverses overdose. Act quickly!

Peel

Peel back the foil and remove the nasal spray. their nostril.

Gently Insert the nozzle into

Place

Press the plunger all the way in until it clicks.

Press





Naloxone nasal spray takes 2-3 minutes to work! it at least 2 minutes for a response before giving more CALL 911! Give rescue breaths while you wait.

A person who has overdosed may wake up after naloxone administration

or they may remain unconscious.

If someone who received naloxone is breathing slowly, shallowly or not at all, rescue breathing is essential.

Give Rescue Breaths!

- Place them on their back. Make sure nothing is their mouth.
 Apply a barrier mask, if available. Tilt their head back, lift their chin and pinch their nose closed. This opens the airway.
- 3. Give one breath slowly, watching to see their chest rise.
- 4. Continue giving one breath every five seconds.
- If they start to gurgle or breathe on their own, stop and roll them onto their side in recovery position.

Steps 1-2 Steps 3-4



A person who has overdosed may wake up after naloxone administration

of they may remain unconscious.

If someone who received naloxone is breathing slowly, shallowly or not at all, rescue breathing is essential

Give Rescue Breaths!

- 1. Place them on their back. Make sure nothing is their mouth.
- Apply a barrier mask, if available. Tilt their head back, lift their chin and pinch their nose closed. This opens the airway.
- 3. Give one breath slowly, watching to see their chest rise.
- 4. Continue giving one breath every five seconds.
- If they start to gurgle or breathe on their own, stop and roll them onto their side in recovery position.

Steps 1-2 Steps 3-4



A person who has overdosed may wake up after naloxone administration

or they may remain unconscious.

If someone who received naloxone is breathing slowly, shallowly or not at all, rescue breathing is essential.

Give Rescue Breaths!

- Place them on their back. Make sure nothing is their mouth.
 Apply a barrier mask, if available. Tilt their head back, lift their chin and pinch their nose closed. This opens the airway.
- 3. Give one breath slowly, watching to see their chest rise.
- 4. Continue giving one breath every five seconds.
- If they start to gurgle or breathe on their own, stop and roll them onto their side in recovery position.

Steps 1-2 Steps 3-4



A person who has overdosed may wake up after naloxone administration

or they may remain unconscious.

If someone who received naloxone is breathing slowly, shallowly or not at all, rescue breathing is essential.

Give Rescue Breaths!

- 1. Place them on their back. Make sure nothing is their mouth.
- Apply a barrier mask, if available. Tilt their head back, lift their chin and pinch their nose closed. This opens the airway.
- 3. Give one breath slowly, watching to see their chest rise.
- 4. Continue giving one breath every five seconds.
- If they start to gurgle or breathe on their own, stop and roll them onto their side in recovery position.

Steps 1-2 Steps 3-4 Step 5

A person who has overdosed may wake up after naloxone administration

of they may remain unconscious.

If someone who received naloxone is breathing slowly, shallowly or not at all, rescue breathing is essential

Give Rescue Breaths!

- Place them on their back. Make sure nothing is their mouth.
 Apply a barrier mask, if available. Tilt their head back, lift
- their chin and pinch their nose closed. This opens the airway
- Give one breath slowly, watching to see their chest rise.

Continue giving one breath every five seconds.

If they start to gurgle or breathe on their own, stop and roll them onto their side in recovery position.

Steps 1-2





A person who has overdosed may wake up after naloxone administration

or they may remain unconscious.

If someone who received naloxone is breathing slowly, shallowly or not at all, rescue breathing is essential

Give Rescue Breaths!

- 1. Place them on their back. Make sure nothing is their mouth.
- Apply a barrier mask, if available. Tilt their head back, lift their chin and pinch their nose closed. This opens the airway.
- Give one breath slowly, watching to see their chest rise.
- 4. Continue giving one breath every five seconds.
- If they start to gurgle or breathe on their own, stop and roll them onto their side in recovery position.

Steps 1-2





A person who has overdosed may wake up after naloxone administration

they may remain unconscious.

If someone who received naloxone is breathing slowly, shallowly or not at all, rescue breathing is essential

Give Rescue Breaths!

- Place them on their back. Make sure nothing is their mouth.
 Apply a barrier mask, if available. Tilt their head back, lift their chin and pinch their nose closed. This opens the airway.
- 3. Give one breath slowly, watching to see their chest rise.
- Continue giving one breath every five seconds.
- If they start to gurgle or breathe on their own, stop and roll them onto their side in recovery position.

Steps 1-2





A person who has overdosed may wake up after naloxone administration

of they may remain unconscious.

If someone who received naloxone is breathing slowly, shallowly or not at all, rescue breathing is essential

Give Rescue Breaths!

- 1. Place them on their back. Make sure nothing is their mouth.
- Apply a barrier mask, if available. Tilt their head back, lift their chin and pinch their nose closed. This opens the airway.
- 3. Give one breath slowly, watching to see their chest rise.
- 4. Continue giving one breath every five seconds.
- If they start to gurgle or breathe on their own, stop and roll them onto their side in recovery position.

Steps 1-2

Steps 3 - 4



A person who has overdosed may wake up after naloxone administration

or they may remain unconscious.

If someone who received naloxone is breathing slowly, shallowly or not at all, rescue breathing is essential.

Give Rescue Breaths!

- Place them on their back. Make sure nothing is their mouth.
 Apply a barrier mask, if available. Tilt their head back, lift their chin and pinch their nose closed. This opens the airway.
- 3. Give one breath slowly, watching to see their chest rise.
- 4. Continue giving one breath every five seconds.
- If they start to gurgle or breathe on their own, stop and roll them onto their side in recovery position.

Steps 1-2





A person who has overdosed may wake up after naloxone administration

or they may remain unconscious.

If someone who received naloxone is breathing slowly, shallowly or not at all, rescue breathing is essential.

Give Rescue Breaths!

- Place them on their back. Make sure nothing is their mouth.
 Apply a barrier mask, if available. Tilt their head back, lift
- their chin and pinch their nose closed. This opens the airway. 3. Give one breath slowly, watching to see their chest rise.
- 4. Continue giving one breath every five seconds.
- If they start to gurgle or breathe on their own, stop and roll them onto their side in recovery position.

Steps 1-2



